

On an identity for the volume integral of the square of a vector field. Remark on a paper by A. M. Stewart

Loyal Durand*

Department of Physics, University of Wisconsin-Madison, Madison, Wisconsin 53706

Gubarev, Stodolsky, and Zakharov¹ have noted the following identity for vector fields that vanish sufficiently rapidly at spatial infinity,

$$\int d^3x \mathbf{A}(\mathbf{x}) \cdot \mathbf{A}(\mathbf{x}) = \frac{1}{4\pi} \int d^3x d^3x' \frac{1}{|\mathbf{x} - \mathbf{x}'|} \left[(\nabla \cdot \mathbf{A}(\mathbf{x})) (\nabla' \cdot \mathbf{A}(\mathbf{x}')) + (\nabla \times \mathbf{A}(\mathbf{x})) \cdot (\nabla' \times \mathbf{A}(\mathbf{x}')) \right], \quad (1)$$

and used it to investigate properties of the vector potential in quantum field theory. Stewart² has shown that the identity is also of general interest in classical electromagnetic theory. It can be used, for example, to derive easily interpreted expressions for the energies in time-dependent electric and magnetic fields.

The existence of this identity is not obvious. In Ref. 1 it is not proven, but it is noted that it follows from the momentum-space relation $(\mathbf{k} \times \tilde{\mathbf{A}})^2 = \mathbf{k}^2 \tilde{\mathbf{A}}^2 - (\mathbf{k} \cdot \tilde{\mathbf{A}})^2$ where $\tilde{\mathbf{A}}(\mathbf{k})$ is the Fourier transform of the vector field. The position-space derivation given by Stewart is based on the Helmholtz decomposition of a three-dimensional vector field into irrotational and solenoidal parts. We give here an alternative position-space derivation that uses only familiar operations starting from the relations

$$\delta^3(\mathbf{x} - \mathbf{x}') = -\frac{1}{4\pi} \nabla^2 \frac{1}{|\mathbf{x} - \mathbf{x}'|} \quad (2)$$

for the Dirac delta function, and

$$\nabla \times (\nabla \times \mathbf{A}) = \nabla(\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A}. \quad (3)$$

Both are familiar to students and are used in the solution of Poisson's equation for the potential of a point charge and the derivation of the wave equation from Maxwell's equations.

By using Eq. (2), we obtain

$$\begin{aligned} \int d^3x \mathbf{A}(\mathbf{x}) \cdot \mathbf{A}(\mathbf{x}) &= \int d^3x d^3x' \mathbf{A}(\mathbf{x}) \cdot \delta^3(\mathbf{x} - \mathbf{x}') \mathbf{A}(\mathbf{x}') \\ &= -\frac{1}{4\pi} \int d^3x d^3x' \mathbf{A}(\mathbf{x}) \cdot \left(\nabla'^2 \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) \mathbf{A}(\mathbf{x}'). \end{aligned} \quad (4)$$

If we integrate by parts over \mathbf{x}' and use the assumed rapid vanishing of $\mathbf{A}(\mathbf{x}')$ for $|\mathbf{x}'| \rightarrow \infty$, we can transfer the action of the derivatives from the factor $1/|\mathbf{x} - \mathbf{x}'|$ to $\mathbf{A}(\mathbf{x}')$ without acquiring extra surface terms. We use the identity (3) for the double curl, make another

partial integration, and rewrite the result as

$$\int d^3x \mathbf{A}(\mathbf{x}) \cdot \mathbf{A}(\mathbf{x}) = -\frac{1}{4\pi} \int d^3x d^3x' \mathbf{A}(\mathbf{x}) \cdot \frac{1}{|\mathbf{x} - \mathbf{x}'|} \nabla'^2 \mathbf{A}(\mathbf{x}') \quad (5a)$$

$$\begin{aligned} &= -\frac{1}{4\pi} \int d^3x d^3x' \mathbf{A}(\mathbf{x}) \cdot \frac{1}{|\mathbf{x} - \mathbf{x}'|} \left[\nabla' (\nabla' \cdot \mathbf{A}(\mathbf{x}')) - \nabla' \times (\nabla' \times \mathbf{A}(\mathbf{x}')) \right] \\ &= \frac{1}{4\pi} \int d^3x d^3x' \left[\left(\mathbf{A}(\mathbf{x}) \cdot \nabla' \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) (\nabla' \cdot \mathbf{A}(\mathbf{x}')) \right. \\ &\quad \left. - \mathbf{A}(\mathbf{x}) \cdot \left(\nabla' \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) \times (\nabla' \times \mathbf{A}(\mathbf{x}')) \right]. \end{aligned} \quad (5b)$$

Now $\nabla'(1/|\mathbf{x} - \mathbf{x}'|) = -\nabla(1/|\mathbf{x} - \mathbf{x}'|)$, so with some rearrangement of the vector products, Eq. (5b) becomes

$$\begin{aligned} \int d^3x \mathbf{A}(\mathbf{x}) \cdot \mathbf{A}(\mathbf{x}) &= -\frac{1}{4\pi} \int d^3x d^3x' \left[\left(\mathbf{A}(\mathbf{x}) \cdot \nabla \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) (\nabla' \cdot \mathbf{A}(\mathbf{x}')) \right. \\ &\quad \left. + \left(\nabla \frac{1}{|\mathbf{x} - \mathbf{x}'|} \right) \times \mathbf{A}(\mathbf{x}) \cdot (\nabla' \times \mathbf{A}(\mathbf{x}')) \right]. \end{aligned} \quad (6)$$

A final partial integration with respect to \mathbf{x} gives the desired identity.

The derivation generalizes easily to give an identity of the same form for the volume integral of the product of two different rapidly decreasing vector fields.

* Electronic address: ldurand@hep.wisc.edu

¹ F. V. Gubarev, L. Stodolsky, and V. I. Zakarov, “On the significance of the vector potential squared,” Phys. Rev. Lett. 86 (11), 2220–2222 (2001).

² A. M. Stewart, “On an identity for the volume integral of the square of a vector field,” Am. J. Phys. (preceding paper).